

Einstein-Podolsky-Rosen (EPR) Correlations and Superluminal Interactions

Luiz Carlos Ryff

*Instituto de Física, Universidade Federal do Rio de Janeiro,
Caixa Postal 68528, 21041-972 Rio de Janeiro, Brazil*

E-mail: ryff@if.ufrj.br

March 20, 2017

Abstract

The possible connection between EPR correlations and superluminal interactions is discussed using simple and palpable arguments. It is shown how an experiment based on time-like events can allow us to answer the question “Can a measurement performed on one of the photons of an entangled pair change the state of the other?” The theorem on superluminal finite-speed causal influences and superluminal signaling is reexamined. It is shown how faster-than-light interactions and Lorentz transformations might peacefully coexist.

1 Introduction

Quantum entanglement, besides being an active topic of research [1], with possible practical consequences in our daily life, has also raised controversial issues related to the foundations of quantum mechanics. The term “EPR correlations” was born from the attempt by Einstein, Podolsky, and Rosen to demonstrate the incompleteness of quantum mechanics using quantum entanglement [2]. Enlightening contributions to the subject have been given by Schrödinger [3] and Bohm [4], and the seminal analysis by Bell [5] of the EPR correlations has led to the notion of quantum nonlocality. Physicists are divided on how to interpret this phenomenon [6], and it is not my intention to discuss the different approaches to the theme. Here, I will start from the assumption that there are only two reasonable explanations for the observed correlations between distant events: a previously shared property or some kind of interaction [7]. Since Bell’s theorem excludes the first alternative, only the second remains, irrespective of how improbable it sounds [8]. This immediately raises the question about the propagation speed of this possible interaction. This problem has been attacked by Gisin’s group. Experiments have been performed to try to determine lower bounds to this speed [9], and it has been shown that if EPR correlations result from superluminal finite-speed causal influences then superluminal signaling is

possible, at least in principle [10]. In this paper I intend to show how simple and palpable (but by no means less rigorous) reasonings help us to clarify some important conceptual questions related to the subject. The paper is organized as follows: In section 2 an argument supporting Bell's point of view according to which EPR correlations strongly suggest that "behind the scenes something is going faster than light" [11] will be introduced; in section 3 the theorem on superluminal finite-speed causal influences and superluminal signaling will be reexamined; in section 4 we will investigate how superluminal interactions and special relativity (strictly speaking, Lorentz transformations) might peacefully coexist. (Please note that I am not saying that EPR correlations mediated by superluminal finite-speed causal influences necessarily lead to superluminal communication) Discussion and conclusion will be presented in section 5.

2 EPR Correlations and Faster-Than-Light (FTL) Interaction

To try to answer the question "Can a measurement performed on one of the photons of an entangled pair change the state of the other?" let us consider the ideal experiment represented in Fig. 1. A source (S) emits pairs of photons (ν_1 and ν_2) that propagate in opposite directions in the polarization-entangled state

$$\begin{aligned} |\psi\rangle &= \frac{1}{\sqrt{2}}(|a_{\parallel}\rangle_1 |a_{\parallel}\rangle_2 + |a_{\perp}\rangle_1 |a_{\perp}\rangle_2) = \frac{1}{\sqrt{2}}(|b_{\parallel}\rangle_1 |b_{\parallel}\rangle_2 + |b_{\perp}\rangle_1 |b_{\perp}\rangle_2) \\ &= \frac{1}{\sqrt{2}}(|c_{\parallel}\rangle_1 |c_{\parallel}\rangle_2 + |c_{\perp}\rangle_1 |c_{\perp}\rangle_2) = \dots, \end{aligned} \quad (1)$$

where the ket $|a_{\parallel}\rangle_1$ ($|a_{\perp}\rangle_1$) represents a photon ν_1 with polarization parallel (perpendicular) to $\mathbf{a}$, and so on [12]. Eq. (1) emphasizes the rotational symmetry of the situation and the fact that the photons have no privileged polarization. A detour has been introduced to have ν_1 always detected before ν_2 in all Lorentz frames. Since the detections of ν_1 and ν_2 are events separated by a time-like interval, there is no doubt that ν_1 is *really* detected first. Therefore, when ν_1 is found in state $|a_{\parallel}\rangle$ ($|a_{\perp}\rangle$), for instance, we immediately know from (1) that ν_2 has been forced into state $|a_{\parallel}\rangle$ ($|a_{\perp}\rangle$). This can be checked using polarizer II and observing that the transmission of ν_2 satisfies Malus' law; or, at least in principle, introducing a third photon (ν_3) identical to ν_2 and performing a HOM experiment [13]. Similarly, if ν_1 is found in state $|b_{\parallel}\rangle$ ($|b_{\perp}\rangle$), ν_2 is forced into state $|b_{\parallel}\rangle$ ($|b_{\perp}\rangle$), and so on. In other words, by "playing" with the orientation of polarizer I it is possible to force ν_2 into states $|a_{\parallel}\rangle$ and $|a_{\perp}\rangle$, or $|b_{\parallel}\rangle$ and $|b_{\perp}\rangle$, or any other pair of mutually orthogonally polarized states. Naturally, if the detection of ν_1 had no influence on ν_2 , there would be no reason for ν_2 to be found only in states $|a_{\parallel}\rangle$ and $|a_{\perp}\rangle$, for instance (or only in states $|b_{\parallel}\rangle$ and $|b_{\perp}\rangle$, and so on). On the other hand, this influence must still be present when space-like events are considered, since the very same correlations can be observed. In an attempt to get rid of this "spooky" action-at-a-distance, we might assume

that photons could be in new states (not contemplated by present physics, however) in which, somehow, different polarizations could coexist simultaneously [14]. However, this solution can be discarded by Bell's theorem [5].

Two points are worth emphasizing in our argument. (a) It is possible to know which photon is really detected first. This is important if we want to investigate if, by acting on a photon of an entangled pair, we can force the other distant photon into a well-defined state. (b) It makes it clear that the second photon is indeed forced into a well-defined polarization state. In fact, if ν_1 is transmitted (reflected) at polarizer I, we know that ν_2 will impinge on polarizer II in state $|a_{\parallel}\rangle$ ($|a_{\perp}\rangle$) (assuming that polarizer I is oriented parallel to $\mathbf{a}$ and the entangled state is represented by (1)), that is, the very same state we would obtain if ν_2 had been transmitted (reflected) after impinging on a polarizer oriented parallel to $\mathbf{a}$. Therefore, *we can predict* the outcome of an experiment involving ν_2 (using polarizer II, for instance, or introducing a third photon, to perform a HOM experiment). Naturally, if we had no information about ν_1 (*which is not the case*), it would be impossible to distinguish a random sequence of photons ν_2 in states $|a_{\parallel}\rangle$ and $|a_{\perp}\rangle$ from another random sequence of photons ν_2 in states $|b_{\parallel}\rangle$ and $|b_{\perp}\rangle$, and so on.

Although we have only discussed an ideal situation, quantum mechanics predictions for EPR correlations have been corroborated by experiment [15]. This strongly suggests that something is indeed going faster than light. At least, this is a possibility that deserves to be investigated.




Fig.1 A source (S) emits a pair of polarization-entangled photons (ν_1 and ν_2) that propagate in opposite directions and impinge respectively on two-channel polarizers (I and II). A detour is introduced to have time-like events in which ν_1 is always detected before ν_2 . Therefore, knowing the state in which ν_1 has been found it is possible to know the polarization state of ν_2 before it impinges on polarizer II. (The arrows emerging from the polarizers indicate the positions of the detectors)

3 The Theorem on Superluminal Finite-Speed Causal Influences

In this section, I will present an alternative and more palpable version of the theorem on superluminal finite-speed causal influences [10]. The assumption of a superluminal causal influence linking space-like separated events raises a problem from the start: how to know which one is the cause and which one is the effect [16]? A way to overcome this limitation (to be discussed in the next section) is introducing a privileged frame in which the “actual” time sequence of the events can be known [17]. In this frame it would be possible to determine which photon is “really” detected first (even in the case of space-like events), forcing the other (or the others) into a well-defined polarization state. Let us then consider the experiment represented in Fig. 2, which we assume as being performed in this hypothetical coordinate system at rest in a Newtonian absolute space [18], and the three-photon polarization-entangled state [19]

$$|GHZ\rangle = \frac{1}{\sqrt{2}}(|H\rangle_1 |H\rangle_2 |H\rangle_3 + |V\rangle_1 |V\rangle_2 |V\rangle_3), \quad (2)$$

where the letters GHZ stand for Greenberger, Horne, and Zeilinger [20], and, taking as a reference the plane on which the photons propagate, the ket $|H\rangle_1$ ($|V\rangle_1$) represents a photon ν_1 with horizontal (vertical) polarization, and so on. Photon ν_1 is sent to Alice (**A**) and ν_2 and ν_3 are sent to Bob (**B**) and Charlie (**C**), who work in the same lab. At instant t_A (in the privileged frame), **A** may decide to measure the polarization state of ν_1 , or not; and, at instant $t_L > t_A$ (also in the privileged frame), **B** and **C** will measure the polarization states of ν_2 and ν_3 . The polarizers are oriented to have the photons emerging either in state $|H\rangle$ (transmitted) or in state $|V\rangle$ (reflected). The condition $\bar{u} > l/(t_L - t_A) > c$ has to be fulfilled, where l is the distance from **A** to **B** and from **A** to **C**, and $\bar{u}$ is the finite superluminal speed. (Although we are considering a very idealized situation, in which a “measurement” occurs at a precise instant and takes no time to be completely accomplished, this will not invalidate the essence of our reasoning.) Supposing that the correlations are purely nonlocal, whenever **B** and **C** perform their measurements, but **A** does not perform hers, the probability of **B** and **C** observing the same outcome is $1/2$, since there can be no communication between them ($\bar{u} < \infty$). On the other hand, whenever **B** and **C** perform their measurements, and **A** performs hers, this probability is equal to 1, since the first measurement forces the other two photons into the same state. Therefore, if we have in the left lab many **As**, and in the right distant lab the corresponding **Bs** and **Cs**, and the **As** combine to take the same decision together, that is, to perform a measurement or not, the **Bs** and **Cs** will know, comparing their results (disregarding improbable statistical fluctuations), what has been decided in the left lab before this information could reach them transmitted by a light signal.

The above discussion may give rise to two important issues. The first has to do with the particular GHZ state (2). It may be argued that if we had a mixture (instead of state (2)) in which $\frac{1}{2}$ of the photons were emitted in the state $|H\rangle_1 |H\rangle_2 |H\rangle_3$ and $\frac{1}{2}$ in the state $|V\rangle_1 |V\rangle_2 |V\rangle_3$, whenever we measured the

horizontal and vertical polarizations, we would obtain the quantum mechanical results, with no need of assuming a nonlocal interaction. However, this would go against the quantum mechanical formalism, which ascribe no polarization to individual photons in state (2). Moreover, as has been shown [19,20], state (2) is an entangled state that, differently from the mixture we are considering, violates locality. Therefore, the correlations involved in the above discussion are indeed nonlocal [21]. (If we had considered the most general situation, in which a fraction p of the photons is emitted in state (2), a fraction $\frac{1-p}{2}$ in state $|H\rangle_1 |H\rangle_2 |H\rangle_3$, and a fraction $\frac{1-p}{2}$ in state $|V\rangle_1 |V\rangle_2 |V\rangle_3$, our argument would not be changed in any essential way.)

The second issue has to do with the duration of a measurement. In principle, the nonlocal connection between Bob and Charlie may be arbitrarily fast. Thus, we can imagine that the time it takes for a measurement to be accomplished is longer than the time necessary for Bob's and Charlie's respective systems to exchange information and to arrive, so to speak, at an agreement on which polarization the photons will be found. We may even assume that the measurement is only concluded when this agreement is reached. In this case, it would be possible to have superluminal finite-speed causal influences without superluminal signaling. Therefore, even considering this imagined possibility as far too unnatural, from a strictly logical point of view it is not immediately obvious that finite-speed superluminal interactions would necessarily lead to the possibility of superluminal communication.




Fig.2 A source (S) emits three polarization-entangled photons (ν_1 , ν_2 , and ν_3) toward three two-channel polarizers, A (for Alice), B (for Bob), and C (for Charlie). Assuming a superluminal finite-speed causal influence, whenever Alice decides to “remove” her polarizer, ν_2 and ν_3 , which are detected at the same time, may be found in different polarization states. (The arrows emerging from the polarizers indicate the positions of the detectors)

4 Superluminal Interaction and Breaking of the Lorentz Symmetry

It might be argued that superluminal finite-speed causal influences should be rejected from the start, since they might lead to FTL communication, and, as a consequence, to causal paradox [16]. Or that they would force us to abandon special relativity altogether. However, things may not be that simple. In the case of the violation of parity, the reflected images of some physical phenomena do not exist in the real world. Similarly, we may assume, *in the case of nonlocal interactions*, that the equivalence between passive and active Lorentz transformations does not exist. In other words, some events that occur in the moving frame cannot take place in the privileged one. More specifically, we will assume that in the latter the speed of the superluminal nonlocal interaction is a constant, irrespective of the velocity of the source or the direction of propagation. Naturally, if Lorentz transformations remain valid, this will not hold in the former (for instance, this interaction will not propagate isotropically, allowing us, in principle, to determine the velocity of the moving frame relative to the privileged frame). Surely, the constancy of the FTL speed prevents the arising of causal paradoxes in the privileged frame, and since events that are coincident in space and time are still coincident under Lorentz transformations, no paradoxes are to be expected in the moving frames; however, it can be instructive to see how things work [22]. Let us consider a pair of reference frames, $\mathbf{S}$ and $\mathbf{S}'$, in the standard configuration, where $\mathbf{S}$ is the privileged frame and $\mathbf{S}'$ moves with velocity $v < c$ along the x axis. Assuming that the Lorentz transformations

$$x' = \gamma(x - vt), \quad (6)$$

$$t' = \gamma\left(t - \frac{v}{c^2}x\right), \quad (7)$$

$$x = \gamma(x' + vt'), \quad (8)$$

and

$$t = \gamma\left(t' + \frac{v}{c^2}x'\right), \quad (9)$$

connect the $\mathbf{S}$ and $\mathbf{S}'$ coordinates (with $\gamma = 1/\sqrt{1 - v^2/c^2}$), we derive

$$u' = \frac{u - v}{1 - \frac{vu}{c^2}} \quad (10)$$

and

$$u = \frac{u' + v}{1 + \frac{vu'}{c^2}} \quad (11)$$

for the velocities.

Let us initially see how the causal paradox arises in special relativity (in which there is no privileged frame and $\mathbf{S}$ and $\mathbf{S}'$ are equivalent). Let the positive quantity $\bar{u} > c$ represent the superluminal signal speed in $\mathbf{S}$. From (10), we see that if $u = \bar{u}$, we can choose v so as to have $v\bar{u}/c^2 > 1$, which leads to $u' < 0$ (with $|u'| > c$ but $\neq \bar{u}$). Therefore, in $\mathbf{S}'$ the signal propagates backwards.

Similarly, from (11) we see that, if $u' = -\bar{u}$, we can choose a v that leads to $u > 0$ (with $u > c$ and $\neq \bar{u}$). That is, in $\mathbf{S}$ the direction of propagation of the signal is reversed. It is this change of direction when we go from $\mathbf{S}$ to $\mathbf{S}'$, and then from $\mathbf{S}'$ to $\mathbf{S}$, that is at the origin of the causal paradox. To see this, let us consider a superluminal signal emitted from $x_0 = 0$, at instant $t_0 = 0$, and reaching $x_1 > 0$ at instant t_1 given by

$$t_1 = \frac{x_1}{\bar{u}} \quad (12)$$

in $\mathbf{S}$. In $\mathbf{S}'$, the signal is transmitted from $x'_0 = 0$, at instant $t'_0 = 0$, reaching x_1 at instant

$$t'_1 = \gamma \left(t_1 - \frac{v}{c^2} x_1 \right) = \gamma \left(1 - \frac{v\bar{u}}{c^2} \right) \frac{x_1}{\bar{u}}, \quad (13)$$

according to (7) and (12). We see that $v\bar{u}/c^2 > 1 \rightarrow t'_1 < 0$. Therefore, in $\mathbf{S}'$ the signal reaches x_1 before it is sent from x_0 (actually, the signal is seen to propagate from x_1 to x_0). But this does not yet represent a paradox, since no contradiction (A and $\neg A$, for instance) is occurring. Let us then determine the point x'_1 in $\mathbf{S}'$ that coincides with x_1 at the instant at which the signal arrives. Using (6) and (12), we obtain

$$x'_1 = \gamma \left(x_1 - v \frac{x_1}{\bar{u}} \right) = \gamma \left(1 - \frac{v}{\bar{u}} \right) x_1. \quad (14)$$

An observer at x'_1 can then send a return signal with $u' = -\bar{u}$ that will take the time of

$$\delta t' = \frac{x'_1}{\bar{u}} = \gamma \left(1 - \frac{v}{\bar{u}} \right) \frac{x_1}{\bar{u}} \quad (15)$$

to arrive at x'_0 . This can lead to a paradox if

$$t'_1 + \delta t' < 0, \quad (16)$$

that is, if the return signal reaches the origin of $\mathbf{S}'$ before t'_0 , namely before the first signal has been sent. This enables an observer in this region, after receiving the return signal, to inform another observer, at the origin of $\mathbf{S}$, not to send the signal. As a consequence, if the signal is sent, it is possible to send a return signal to impede the emission of the signal. That is, the signal would be sent and not sent at the same time! Let us see the condition v would have to fulfil. From (16), (15), and (13), we obtain

$$\gamma \left(1 - \frac{v\bar{u}}{c^2} \right) \frac{x_1}{\bar{u}} + \gamma \left(1 - \frac{v}{\bar{u}} \right) \frac{x_1}{\bar{u}} < 0, \quad (17)$$

which leads to

$$v > \frac{2\bar{u}}{1 + \frac{\bar{u}^2}{c^2}}. \quad (18)$$

Since the right-hand side of (18) is always smaller than c , it is always possible to find a v that satisfies the above condition; therefore, we would indeed have a paradox.

Now let us see how the existence of a privileged frame in which the superluminal speed is a constant does not lead to a causal paradox. Instead of (15), we have

$$\delta t' = \frac{x_1'}{-\bar{u}'} = -\gamma \left(1 - \frac{v}{\bar{u}}\right) \frac{x_1}{\bar{u}}, \quad (19)$$

where the velocity of the return signal (using (10)) is

$$\bar{u}' = \frac{-\bar{u} - v}{1 + \frac{v\bar{u}}{c^2}}. \quad (20)$$

The condition to have a causal paradox is then

$$\gamma \left(1 - \frac{v\bar{u}}{c^2}\right) \frac{x_1}{\bar{u}} - \gamma \left(1 - \frac{v}{\bar{u}}\right) \frac{x_1}{\bar{u}'} < 0, \quad (21)$$

instead of (17), where (19), (16), and (13) have been used. From (20) and (21) we obtain

$$v > c, \quad (22)$$

which contradicts our initial assumption that the velocity of reference frame $\mathbf{S}'$ is slower than the velocity of light. As a consequence, there can be no causal paradox.

5 Discussion

As we have seen, quantum mechanical formalism strongly suggests that “measuring” the polarization of a photon of a polarization-entangled pair [23] forces the other into a well-defined polarization state. This is particularly apparent when we consider time-like events. Moreover, since the same correlations are predicted when space-like events are taken into account, this seems to imply some kind of superluminal interaction between the photons and, as a consequence, the existence of a privileged frame in which the “real” time order of the physical events could be known. Furthermore, if the speed of this FTL interaction is finite, then superluminal signaling becomes possible in principle. However, as has been shown, assuming that the Lorentz symmetry is broken in the case of superluminal interactions (and only in this case), this does not lead to causal paradoxes or to the abandonment of the well-established laws of physics which are Lorentz covariant. Therefore, the possible connection between EPR correlations and FTL interactions is a subject that deserves to be investigated. In my opinion, we should keep in mind Poincaré’s wise warning: “... le principe de relativité physique est un fait expérimental, au même titre que les propriétés des solides naturels; comme tel, il est susceptible d’une incessante révision...” [24] (the principle of physical relativity is an experimental fact, like the properties of natural solids; as such, it is susceptible to incessant revision) [25].

References

- [1] Bub, J.: <http://plato.stanford.edu/archives/win2010/entries/qt-entangle/>; Tittel, W., Gregor Weihs, G.: Quantum Information and Computation **1**, 3 (2001); Bouwmeester, D., Ekert, A., Zeilinger, A. (eds.): The Physics of Quantum Information. Springer (2000); Gibney, E.: “One giant step for quantum internet”, in Nature **535**, 478 (2016)
- [2] Einstein, A., Podolsky, B., Rosen, N.: Phys. Rev. **47**, 777 (1935). Reprinted in: Wheeler, J.A., Zurek, W.H. (eds.): Quantum Theory and Measurement. Princeton University Press, Princeton (1983)
- [3] Schrödinger, E.: Naturwissenschaften **23**, 807, 823, and 844 (1935). English translation originally published in Proceedings of the American Philosophical Society **124**, 323 (1980). Reprinted in: Wheeler, J.A., Zurek, W.H. (eds.): Quantum Theory and Measurement. Princeton University Press, Princeton (1983)
- [4] Bohm, D.: Quantum Theory. Prentice Hall, New York (1951)
- [5] Bell, J.S.: Physics **1**, 195 (1964). Reprinted in: Bell, J.S.: Speakable and Unsayable in Quantum Mechanics. Cambridge University Press, Cambridge (1989) and in Wheeler, J.A., Zurek, W.H. (eds.): Quantum Theory and Measurement. Princeton University Press, Princeton (1983)
- [6] To have an idea on how divided are physicists on foundational questions on quantum mechanics see Schlosshauer, M. (ed.): Elegance and Enigma. Springer, Berlin (2011), and also Schlosshauer, M., Kofler, J., Zeilinger, A.: arXiv: 1301.1069v1 [quant-ph], Sommer, C.: arXiv: 1303.2719v1 [quant-ph], and Norsen, T., Nelson, S.: arXiv: 1306.4646v2 [quant-ph]
- [7] Naturally, discarding very improbable coincidences. Ryff, L.C.: arXiv: 1308.5690v1 [quant-ph]. The reason for this (disputable?) attitude will become evident in section 2. Actually, there are different (and conflicting) points of view on Bell’s inequalities and quantum nonlocality. For the interested reader I recommend Bell, M. and Gao, S. (ed.): Quantum Mechanics and Reality: 50 Years of Bell’s Theorem, Cambridge University Press, Cambridge (2016)
- [8] “Eliminate all other factors, and the one which remains must be the truth.” “How often have I said to you that when you have eliminated the impossible, whatever remains, however improbable, must be the truth?” (Sherlock Holmes - Wikiquote).
- [9] Scarani, V., Tittel, W., Zbinden, H., Gisin, N.: Phys. Lett. A **276**, 1 (2000); Salart, D., Baas, A., Branciard, C., Gisin, N., Zbinden, H.: Nature **454**, 861 (2008). See also Yin, J. et al.: Phys. Rev. Lett. **110**, 260407 (2013)

- [10] Scarani, V., Gisin, N.: Phys. Lett. A **295**, 167 (2002); Scarani, V., Gisin, N.: Braz. J. Phys. **35**, 328 (2005); Bancal, J.-D., et al.: Nat. Phys. **8**, 867 (2012); Barnea, T.J., et al.: Phys. Rev. A **88**, 022123 (2013)
- [11] Bell, J.S., in an interview in Davies, P.C.W., Brown, J.R. (eds.): The Ghost in the Atom. Cambridge University Press, Cambridge (1989). In the same book, Bohm explicitly entertains the idea of superluminal signaling
- [12] Kwiat, P.G., Waks, E., White, A.G., Appelbaum, I., Eberhard, P.H.: Phys. Rev. A **60**, R773 (1999)
- [13] Hong, C. K., Ou, Z. Y., Mandel, L.: Phys. Rev. Lett. **59**, 2044 (1987)
- [14] Let us imagine a situation in which the polarizers in Fig. 1 have the same orientation. According to (1), whenever ν_1 is transmitted (reflected), ν_2 must also be transmitted (reflected). Therefore, assuming locality (i.e., what happens to ν_1 cannot affect ν_2 , and vice versa), whether ν_1 and ν_2 will be transmitted or reflected is already determined before they impinge on the polarizers; otherwise, we could have a situation in which one photon is transmitted and the other is reflected. In other words, perfect correlations and locality imply a strong form of determinism. Assuming that the source has no information about the orientations of the polarizers, each photon pair has to be emitted with “instructions,” so to speak, for all possible orientations. For instance, transmission in case of orientation **a**, reflection in case of orientation **b**, and so on. Further details can be found in Ryff, L. C.: arXiv: 0510101[quant-ph]
- [15] Clauser, J. F., Shimony, A.: Rep. Prog. Phys. **41**, 1881 (1978); Aspect, A.: Thèse d’Etat (Orsay, 1983); Shi, Y., H., Alley, C., O.: Phys. Rev. Lett. **61**, 2921 (1998); Weihs, G., Jennewein, T., Simon, C., Weinfurter, H., Zeilinger, A.: Phys. Rev. Lett. **81**, 5039 (1998); Pittman T., B., Franson, J., D.: Phys. Rev. Lett. **90**, 240401 (2003)
- [16] Bohm, D.: The Special Theory of Relativity. Benjamin (1965)
- [17] The introduction of a privileged frame is discussed by John Bell in “How to teach special relativity”, reprinted in Bell, J.S.: Speakable and Unspeakable in Quantum Mechanics. Cambridge University Press, Cambridge (1989), and advocated in Bohm, D., Hiley, J. B.: The Undivided Universe: An Ontological Interpretation of Quantum Theory. Routledge, London (1993)
- [18] Our argument is based on the reasoning introduced in Ryff, L.C.: arXiv: 0903.1076 [quant-ph]
- [19] Pan, J.-W., Bouwmeester, D., Daniell, M., Weinfurter, H., Zeilinger, A.: Nature **403**, 515 (2000); Hamel, D., R., et al: arXiv: 1404.7131 [quant-ph]; Pan, J.-W., Zeilinger, A.: “Multi-Photon Entanglement and Quantum Nonlocality”, in Pan, J.-W., Zeilinger, A. (eds.): Quantum [Un]speakables. Springer (2002); Krenn, M., Hochrainer, A., Lahiri, M., Zeilinger, A.: arXiv:1610.00642 [quant-ph], and references there in

- [20] Greenberger, D. M., Horne, M. A., Shimony, A., Zeilinger, A.: Am. J. Phys. **58**, 1131 (1990); Ryff, L. C.: Am. J. Phys. **65**, 1197 (1997)
- [21] Entangled states have been used to suggest seemingly influences from the future into the past (Peres, A.: J. Mod. Opt. **47**, 139 (2000); Jennewein, T., Weihs, G., Pan, J-W., Zeilinger, A.: Phys. Rev. Lett. **88**, 017903 (2002)), however, as it has been shown (Ryff, L. C.: J. Mod. Opt. **48**, 905 (2001); Ryff, L. C.: arXiv: 0303082 [quant-ph]), such inference is unjustified. The interested reader can also see: Jennewein, T., Weihs, G., Pan, J-W., Zeilinger, A.: arXiv: 0303104 [quant-ph], for a reply.
- [22] Ryff, L. C.: arXiv: 1005.5092 [quant-ph]
- [23] Here, the term “measuring” may sound misleading, since, according to the quantum mechanical formalism, before it impinges on the polarizer and is detected, no well-defined polarization can be assigned to the photon. The measurement only “actualizes” a potentiality, so to speak. This does not force us to abandon realism, but only a sort of naive one.
- [24] Poincaré, H.: Dernières Pensées, Chapitre II: L’espace et le temps, in Oeuvres de Henri Poincaré. Editions La Bibliothèque Digitale. Kindle edition (2012)
- [25] Experiments to investigate the possible violation of Lorentz symmetry in the case of entangled particles have been suggested in Ryff, L. C.: arXiv: 0809.0530 [quant-ph] and Ryff, L. C.: Found. Phys. **44** (1), 58 (2014). On the other hand, the connection between EPR correlations, quantum teleportation, and entanglement swapping *via Lorentz transformations* is examined in Ryff, L. C.: Phys. Rev. A **60**, 5083 (1999)